

Ethics in Psychiatry



Ethical Considerations When Treating Patients with Schizophrenia

by Edmund Howe, MD, JD

INTRODUCTION

Within medicine, some values have changed profoundly over the last few decades. Chief among these is truthfulness—specifically, whether physicians should tell patients the truth when they have a potentially fatal illness, such as cancer. Just decades ago, it was

thought that they should not. Now it is thought that they should. Some physicians presently even advocate using advance directives to predetermine whether patients will want to know the truth at a later time.^{1,2} The extent to which this value of truthfulness is now given priority is reflected by the

current, widespread, ethical consensus that generally doctors should tell patients the truth *even when this may cause foreseeable, additional harm to patients*.

In this commentary, I shall discuss five contexts in which the value of respecting the autonomy of patients with schizophrenia may be in conflict with the values of doing good or avoiding harm. The former is a value not based on the patient's outcome or consequences; the latter two are based on consequences. I suggest that psychiatrists seek to balance these different kinds of values so that in some cases they give greater priority to respecting these patients' autonomy, but in other cases give greater priority to trying to benefit these patients medically as much as they can.

1. TELLING HALF-TRUTHS TO PATIENTS WITH SCHIZOPHRENIA WHO ARE PARANOID

The conflict between truth telling and doing good may be faced by any psychiatrist who considers withholding information from a patient regarding his or her likely psychodynamics or utmost underlying emotional needs. Psychiatrists routinely choose to initially withhold certain types of knowledge because if they disclose everything right away, they run the risk of patients never returning and therefore not receiving adequate care. Ethically, the degree to which a psychiatrist withholds information may disrespect a patient's autonomy and may even be regarded as lying by omission. However, this ethical "price" may be warranted due to other competing and mutually exclusive values that may benefit the patient in other ways.

This same conflict may exist when psychiatrists treat patients

with schizophrenia who have paranoia. Most psychiatrists generally believe that to be maximally effective, they should not directly confront the delusions of patients. Andreason and Black state in their recent basic psychiatric textbook, “A patient with delusional disorder may be more accepting of

Empirical testing of Havens’s approach may not even be possible. Havens says that for his approach to be successful, psychiatrists must genuinely believe that their patients’ symptoms are strengths.^{4,5} If he is right, this genuineness may be impossible experimentally to discern!

One patient I saw...reported that she felt emotionally attached to inanimate objects. I knew from prior meetings with her that she had extremely warm relationships with all her adult children. “Of course!” I said. “Your exceptional capacity for caring has, I am sure, spilled over!”

medication if *it is explained as a treatment for the anxiety, dysphoria, and stress that the patient [has due to] his or her delusions.*”³ [Emphasis added] The use of this partial truth may reduce the risk of avoidable harm, but others have carried this same approach still further. Havens uses a related approach to help establish and then enhance the therapeutic alliance.⁴ He urges psychiatrists to identify ways in which these patients’ symptoms are strengths and to then share this view with them.^{4,5} “[O]ne needs,” he says, “to convey respect while inevitably feeling with another part of one’s mind the emptiness of the patient’s claim.”⁴ He explains, “The widespread concentration on pathology, especially before an alliance is cemented, is not only demoralizing, but it deprives the clinician of an ally.”⁴ According to Havens, a clinician might say in response to a patient’s sharing a grandiose thought, for instance, “How wonderful!”⁵

The gains from psychiatrists sharing partial truths may not be evidence-based.^{6,7} Havens states, anecdotally, “Clinicians’ fears will be relieved by [a positive result].”⁴

Though such approaches may not be empirically proven, psychiatrists still may find them useful. One patient I saw, for example, reported that she felt emotionally attached to inanimate objects. I knew from prior meetings with her that she had extremely warm relationships with all her adult children. “Of course!” I said. “Your exceptional capacity for caring has, I am sure, spilled over!”

Forming a relationship by using such half-truths, despite these possibly representing, ethically, lies, is in Havens’s view the most critical first step in treating patients with schizophrenia.⁴ These patients, he asserts, are more likely to feel isolated due to their symptoms, and thus are more likely to be alone.⁴ If a psychiatrist reframes how a patient regards his or her heightened paranoid thoughts so that he or she sees these thoughts as possible strengths, this may enable the psychiatrist to build a stronger patient alliance, which may in turn help the patient feel less isolated and alone.

2. SUPPORTING THE “RISKY” AMBITIONS OF PATIENTS WITH SCHIZOPHRENIA

An emerging shift in both clinical thinking and the underlying ethics of

psychiatry is placing greater emphasis on the quality of life for patients with schizophrenia, as opposed to primarily trying only to give them relief from their symptoms.^{8–11} One way to improve a patient’s quality of life is by allowing more input by the patient regarding his or her care.

Some patients with schizophrenia are exceptionally ambitious. This may cause some conflict for them and for their psychiatrists. A patient with schizophrenia may wish to pursue a competitive career, for example, that seems to the psychiatrist to be beyond the patient’s capabilities due to certain limitations (e.g., intellectual capabilities), as suggested by what the patient has been able to do in the past. To pursue this wish, the patient runs the risk of exacerbating the symptoms of schizophrenia; yet to not do so may diminish the quality of his or her life by not pursuing a lifelong dream. The psychiatrist faces a dilemma when treating such a patient. Does he or she support the patient’s autonomous wish, which may add value to the patient’s life? Or does the psychiatrist counsel the patient against pursuing such an ambitious task, so as to avoid the risk of exacerbating the illness? Perhaps the psychiatrist should be less paternalistic in a case like this by giving more moral weight to what the patient wants to attempt. Even though the psychiatrist may anticipate that the ambitious pursuit of a particular goal by the patient may possibly make the patient’s schizophrenia worse, it may be that the psychiatrist, notwithstanding this, should support the patient in taking this greater risk in support of the patient’s hope that this will result in him or her having an improved quality of life.

Shared decision making^{13,14} is an approach some psychiatrists use with patients with schizophrenia. Using this approach, the

psychiatrist provides the patient with more information and involves him or her more in treatment decisions. This approach, according to Hamann,¹³ “explicitly goes beyond informed consent.” It aims to decrease “the informational and power asymmetry between doctors and patients by increasing the patient’s information and control over treatment decisions.”¹³ This may involve the use of directional aids. These aids may depict for the patient the relative pros and cons of different scenarios (e.g., switching to a different antipsychotic drug), and then the choice is made to a greater extent by the patient.¹³

Respecting patients by respecting their autonomy is a value independent of actual consequences. It may be that this value should prevail, even over a psychiatrist’s more traditional value of protecting patients, despite the fact that patients with schizophrenia may be more prone to losing decision-making capacity.

3. ASKING A PATIENT WITH SCHIZOPHRENIA WHETHER HE OR SHE WANTS TO WRITE PSYCHIATRIC ADVANCE DIRECTIVES

I have mentioned advance directives in the context of truth-telling. Analogously, a psychiatrist can ask a patient with schizophrenia who still has decisional capacity what his or her wishes are regarding his or her care should he or she lose decisional capabilities at a later time. These psychiatric advance directives enable the psychiatrist to pre-determine several future outcomes, including where the patient will be hospitalized, which psychiatrist will treat the patient, and what medications will be given to the patient.^{15–21} By allowing the patient

to participate in these decisions, the psychiatrist is respectful of the patient’s autonomy and also able to provide optimal medical care. Some patients may not want to pursue this option.^{13,22} Something to further consider is that a psychiatrist who presents this option to a patient may cause the patient exceptional fear, which could make the patient’s schizophrenic illness worse.

4. PERSUADING PATIENTS WITH SCHIZOPHRENIA TO INVOLVE THEIR FAMILY MEMBERS

A substantial advance in the treatment of schizophrenia has been the understanding that psychiatrists can benefit patients with schizophrenia by educating their family members about the harmful effects of criticism and negative emotions, especially when they are expressed in an exaggerated way.^{23–27} Family members can further assist a patient with schizophrenia by

withholding of information may be regarded as lying by omission.²² Psychiatrists should not deny to themselves that (some degree of) coercing (or lying) is a genuine risk of making certain interventions. Psychiatrists should know that no action, including being coercive or lying, should ever be ethically prohibited on this basis alone without also considering whether there are other competing moral values in a given case that may be more important. Thus, it may be ethically justifiable for psychiatrists to accept even these actions and other (relative) harms in some instances.

To help illustrate the possible justification of using persuasion, psychiatrists may have an ethical obligation to try to persuade a patient with schizophrenia to continue taking his or her antipsychotic medication, even after only one psychotic episode. This is

Just as it may be ethically justifiable to encourage a patient with schizophrenia to continue taking his or her medication, a psychiatrist may also have some ethical obligation to try to persuade a patient with schizophrenia to involve his or her family more in his or her care. This may be considered coercive, but in the end, it will probably help the patient have a better quality of life.

helping the patient to detect, earlier on, when he or she is “running into trouble,” which may help to avoid hospitalization. Moreover, in regard to the importance of quality of life, a patient’s family members may contribute to the patient’s sense of meaning and joy most uniquely.^{28,29} Thus, it may be beneficial if the psychiatrist can persuade a patient to have his or her family maximally involved in his or her care.

Ethically, attempts at persuasion may be regarded as coercion, just as

because the likelihood of relapse if the medication is stopped is much higher.^{30,31} The losses patients with schizophrenia may suffer due to relapse may be greatest among those who are doing best on medication. This is because they have “more to lose.” Accordingly, a psychiatrist’s ethical obligation to try to persuade such a patient with schizophrenia to continue taking his or her medications may be increased.

Just as it may be ethically justifiable to encourage a patient

with schizophrenia to continue taking his or her medication, a psychiatrist may also have some ethical obligation to try to persuade a patient with schizophrenia to involve his or her family more in his or her care. This may be considered coercive, but in the end, it will probably help the patient have a better quality of life.

5. PRODRAMAL SCHIZOPHRENIA

Perhaps the most difficult ethical question raised in regard to the treatment of patients with schizophrenia over the last decade is if or when a psychiatrist should tell a patient that he or she is at an

increased risk of developing schizophrenia. Suggestive symptoms may also include disturbances of perception and thought, suspiciousness, and grandiosity.³⁷

It is still uncertain which persons with prodromal findings will actually progress to schizophrenia, and less than half of these people actually do.^{35–39} Different prodromal features may have different rates of conversion, but which features develop at what rates is not fully known.

Some psychiatrists believe that telling persons that they are at risk for developing schizophrenia, much

less treating them, is not necessary. “Risk for schizophrenia is generally not mentioned to either patients or family members, since we don’t believe the available information justifies such use of diagnostic labels with only attenuated symptoms.”⁴⁰

Others feel this same way because they believe that far too many of these persons would later turn out to be “false positives” for schizophrenia. Disclosing the full truth may scare these patients and their families profoundly. In addition, some psychiatrists feel that such knowledge would create exceptionally harmful stigma.^{40,41} The stress from being scared and from the stigma may actually “fuel” the development of schizophrenia if a patient is at all predisposed.³⁷

Perhaps the psychiatrist could use an ethical “sliding scale.” If he or she uses this approach, he or she could vary what to say and do with each individual patient, depending on additional prognostic factors...[e.g.,] a patient’s insights, the strength of a patient’s relationships with family members (or others who would support the patient), and the strength of the relationship the patient has with his or her psychiatrist.

Informing a patient of his or her prodromal state may become a self-fulfilling prophecy.

Still other psychiatrists believe that the risks of not treating a patient with prodromal symptoms of schizophrenia are potentially profound, and that whatever can be done to keep the schizophrenia from emerging should be done.⁴² Efforts at preventing schizophrenia could include, for example, attempts to reduce the sources of stress patients experience in their external environments, psychotherapy, use of antidepressants or anti-anxiety drugs, and the prophylactic use of antipsychotic medications.

Still others favor using what might be considered a compromise solution. They may encourage a patient to reduce environmental stresses as much as possible and/or to use antidepressants or anti-anxiety medications.⁴³

Here, an additional ethical question to consider is what effect would telling or not telling a patient that he or she may develop schizophrenia have on the patient/psychiatrist relationship. This would be the question that Havens would give paramount importance.

Perhaps the psychiatrist could use an ethical “sliding scale.” If he or she uses this approach, he or she could vary what to say and do with each individual patient, depending on additional prognostic factors. These factors might include a patient’s insight, the strength of a patient’s relationships with family members (or others who would support the patient), and the strength of the relationship the patient has with his or her psychiatrist. When using this approach, however, the psychiatrist should be mindful not to make idiosyncratic judgments, which may be ethically problematic.

CONCLUSION

Psychiatrists treating patients with schizophrenia may face ethical conflicts. These conflicts often are between helping these patients maximally and respecting their autonomy optimally.

The current consensus is that the value of respecting a patient's autonomy generally should prevail, even when a patient may potentially do worse medically.⁴⁴ Psychiatrists, however, should always carefully consider what moral weight should be given to the values of doing good and avoiding harm, which were given more weight in the past when psychiatrists generally were more paternalistic. This should particularly be the case when they treat patients with schizophrenia since these patients' decision-making capacities are likely to be impaired.

REFERENCES

- Azu MC, Jean S, Piotrowski JM, O'Hea B. Effective methods for disclosing breast cancer diagnosis. *Am J Surg* 2007;194:488–90.
- Keating DT, Nayeem K, Gilmartin JJ, O'Keefe ST. Advance directives for truth disclosure. *Chest* 2005;128:1037–9.
- Andreasen NC, Black DW. *Introductory Textbook of Psychiatry, Third Edition*. Washington, DC: American Psychiatric Publishing, Inc., 2001: 258.
- Havens LL. The best kept secret: How to form an effective alliance. *Harv Rev Psychiatry* 2004;12:56–62.
- Havens LL, Ghaemi SN. Existential despair and bipolar disorder: The therapeutic alliance as a mood stabilizer. *Am J Psychother* 2005;59(2):137–47.
- Stanard RP. The effect of training in a strengths model of case management on client outcomes in a community mental health center. *Community Ment Health J* 1999;35:169–79.
- Weiden PJ. Understanding and addressing adherence issues in schizophrenia: From theory to practice. *J Clin Psychiatry* 2007;68 (suppl 14):14–9.
- Kane JM. Treatment strategies to prevent relapse and encourage remission. *J Clin Psychiatry* 2007;68 (suppl 14):27–30.
- Marwaha S, Johnson S, Bebbington P, et al. Correlates of subjective quality of life in people with schizophrenia: findings from the EuroSC study. *J Nerv Ment Dis* 2008;196:87–94.
- Nordt C, Müller B, Rössler W, Lauber C. Predictors and course of vocational status, income, and quality of life in people with severe mental illness: A naturalistic study. *Soc Sci Med* 2007;65:1420–9.
- MacBeth A, Gumley A. Premorbid adjustment, symptom development and quality of life in first episode psychosis: A systematic review and critical reappraisal. *Acta Psychiatrica Scand* 2008;117:85–99.
- McEvoy JP, Lieberman JA, Stroup TS, et al. Effectiveness of clozapine versus olanzapine, quetiapine, and risperidone in patients with chronic schizophrenia who do not respond to prior psychiatric treatment. *Am J Psychiatry* 2006;163:600–10.
- Hamann J, Langer B, Winkler V, et al. Shared decision making for in-patients with schizophrenia. *Acta Psychiatr Scand* 2006;114:263–73.
- Hudson TJ, Owen RR, Thrush CR, et al. Guideline implementation and patient-tailoring strategies to improve medication adherence for schizophrenia. *J Clin Psychiatry* 2008;69:74–80.
- Elbogen EB, Swartz MS, Van Dorn R, et al. Clinical decision making and views about psychiatric advance directives. *Psychiatr Serv* 2006;57:350–5.
- Henderson C, Swanson JW, Szmukler, et al. A typology of advance statements in mental health care. *Psychiatr Serv* 2008;59(1):63–71.
- Srebnik DS, Rutherford LT, Peto T, et al. The content and clinical utility of psychiatric advance directives. *Psychiatr Serv* 2005;56:592–8.
- Van Citters AD, Naidoo U, Foti ME. Using a hypothetical scenario to inform psychiatric advance directives. *Psychiatr Serv* 2007;58:1467–71.
- Swanson JW, Swartz MS, Elbogen EB, et al. Facilitated psychiatric advance directives: A randomized trial of an intervention to foster advance treatment planning among persons with severe mental illness. *Am J Psychiatry* 2006;163:1943–51.
- Gremmen I, Widdenshoven G, Beekman A, et al. Ulysses arrangements in psychiatry: A matter of good care? *J Med Ethics* 2008;34:77–80.
- O'Reilly RL. The capacity to execute an advance directive for psychiatric treatment. *Int J Law Psychiatry* 2008;31:66–71.
- Widdenshoven G, Berghmans R. Coercion and pressure in psychiatry: lessons from Ulysses. *J Med Ethics* 2007;33:560–3.
- Pickett-Schenk SA, Lippincott RC, Bennett C, Steigman PJ. Improving knowledge about mental illness through family-led education: The journey of hope. *Psychiatr Serv* 2008;59(1):49–56.
- Morken G, Grawe RW, Widen JH. Effects of integrated treatment

- on antipsychotic medication adherence in a randomized trial in recent-onset schizophrenia. *J Clin Psychiatry* 2007;68:566–71.
25. Brent BK, Guiliano AJ. Psychotic-spectrum illness and family-based treatments: A case-based illustration of the underuse of family interventions. *Harv Rev Psychiatry* 2007;15:161–8.
 26. McFarlane WR, Cook WL. Family expressed emotion prior to onset of psychosis. *Fam Process* 2007;46:185–97.
 27. Cohen AN, Glynn SM, Murray-Swank, et al. The family forum: Directions for the implementation of family psychoeducation for severe mental illness. *Psychiatr Serv* 2008;59:40–8.
 28. Greenberg JS, Knudsen KJ, Aschbrenner KA. Prosocial family processes and the quality of life of persons with schizophrenia. *Psychiatr Serv* 2006;57:1771–7.
 29. Smith MJ, Greenberg JS. Factors contributing to the quality of sibling relationships for adults with schizophrenia. *Psychiatr Serv* 2008;59:57–62.
 30. Robinson D, Woerner MG, Alvir JM, et al. Predictors of relapse following response from a first episode of schizophrenia or schizoaffective disorder. *Arch Gen Psychiatry* 1999;56:241–7.
 31. Law MR, Soumerai, Ross-Degnan D, Adams AS. A longitudinal study of medication nonadherence and hospitalization risk in schizophrenia. *J Clin Psychiatry* 2008;69:47–53.
 32. Filakovic P, Degmele D, Kole E, Benic D. Ethics of the early intervention in the treatment of schizophrenia. *Psychiatria Danubina* 2007;19(3):209–15.
 33. Cannon TD, Cadenhead K, Cornblatt B, et al. Prediction of psychosis in youth at high clinical risk: a multisite longitudinal study in North America. *Arch Gen Psychiatry* 2008;65:28–37.
 34. McGorry PD, Yung AR, Bechdolf A, Amminger P. Back to the future: Predicting and reshaping the course of psychotic disorder. *Arch Gen Psychiatry* 2008;65:25–7.
 35. Addington J, Cadenhead KS, Cannon TD, et al. North American prodrome longitudinal study: A collaborative multisite approach to prodromal schizophrenia research. *Schizophr Bull* 2007;33:665–72.
 36. Schultze-Lutter F, Ruhrmann S, Hoyer C, et al. The Initial prodrome of schizophrenia: Different duration, different underlying deficits? *Compr Psychiatry* 2007;48:479–88.
 37. Corcoran C, Malaspina D, Hercher L. Prodromal interventions for schizophrenia vulnerability: The risks of being “at risk.” *Schizophr Res* 2005;73:173–84.
 38. Cannon TD, Cornblatt B, McGorry P. The empirical status of the ultra high-risk (prodromal) research paradigm. *Schizophr Bull* 2007;33:661–4.
 39. Yung AR, Yuen HP, Berger G, et al. Declining transition rate in ultra high risk prodromal services: Dilution or reduction of risk? *Schizophr Bull* 2007;33:673–81.
 40. Cornblatt BA, Lencz T, Kane JM. Treatment of the schizophrenia prodrome: Is it presently ethical? *Schizophr Res* 2001;51:31–8.
 41. Day JC, Bentall RP, Roberts C, et al. Attitudes toward antipsychotic medication. *Arch Gen Psychiatry* (2005);62:717–24.
 42. McGlashan TH, Addington J, Cannon, et al. Recruitment and treatment practices for help-seeking “prodromal” patients. *Schizophr Bull* 2007;33:715–26.
 43. Cornblatt BA, Lencz T, Smith CW, et al. Can antidepressants be used to treat the schizophrenia prodrome? Results of a prospective, naturalistic treatment study of adolescents. *J Clin Psychiatry* 2007;68(4):546–7.
 44. Boyle RJ. Communications, truth-telling, and disclosure. In: Fletcher JC, Spencer EM, Lombardo PA. *Fletcher's Introduction to Clinical Ethics, Third Edition*. Hagerstown, MD: University Publishing Group, Inc., 2005. ●

AUTHOR AFFILIATION: Dr. Howe is Professor, Department of Psychiatry, and Director, Programs in Medical Ethics, Senior Scientist, Center for the Study of Traumatic Stress, Uniformed Services University of the Health Sciences, Bethesda, Maryland.

ADDRESS CORRESPONDENCE TO: Edmund Howe, MD, JD, A1040A, USUHS, 4301 Jones Bridge Rd., Bethesda, MD 20814-4799; E-mail: ehowe@usuhs.mil